


# Synthetic LSD / N-Bomb Information Sheet

The chemicals are distributed onto perforated sheets of blotter paper, commonly covered in colorful designs or images.




Squares are torn off of the perforated sheets into individual portions, or "hit." These squares may contain a whole or partial image.



Squares can be ingested orally or chemicals can be absorbed through the skin.



The chemicals are commonly stored and transported in aluminum foil or small plastic bags to avoid contact with the skin.



# ***Synthetic LSD / N-Bomb Information Sheet***

## ***Facts about 25I-NBOMe (known as 25I)***

**First developed in 2003, but did not become highly recreational until 2010.  
Initially sold by companies specializing in “research chemicals.”**

- **Can be used:**
  - Orally – (eaten, ingested)
  - Sublingual – (tabs held in the mouth)
  - Buccal – (tabs held in the mouth)
  - Nasal/intranasal – (nasal spray)
  - IV injection
  - Smoking in powdered form
- **Effects usually last 6-10 hours, depending on dosage**
- **Similar effects as LSD:**
  - **Perceived positive effect:**
    - Strong open and closed eye visuals, including trails and color shifts
    - Mood lift
    - Euphoria
    - Mental and physical stimulation
    - Increase in associative and creative thinking
    - Increased awareness and appreciation of music
    - Life-changing spiritual experiences
    - Feeling of love and empathy
  - **Negative effects:**
    - Tachycardia – (increased heart rate)
    - Hypertension – (elevated blood pressure)
    - Agitation, confusion and difficulty focusing
    - Aggression, visual and auditory hallucinations
    - Seizures
    - Hyperpyrexia – (very high fever)
    - Clonus – (involuntary, rhythmic, muscular contractions and relaxations)
    - Elevated white blood cell count
    - Elevated creatine kinase – (seen in heart attacks, when the heart muscle is damaged, or in conditions that produce damage to the skeletal muscles or brain)
    - Acute kidney injury
    - Death
  - **Observing someone under the influence:**
    - Confusion
    - Scrambled communication
    - Tunnel vision
    - Vasoconstriction – (narrowing of blood vessels and veins – causes cold, clammy skin)
    - Nausea and vomiting
    - Paranoia, fear, and panic
    - Shaking